

Original Article

Systematic Review: Safety and Efficacy Comparison of Laser Therapy With And Other Resurfacing Technology

Hettihewa L.M

Department of Health and Medical Science Faculty of Health Sciences, CINEC Campus

Menik.Hettihewa@cinec.edu

ABSTRACT

There is an increasing demand of almost all young and adult population for the noninvasive cosmetic procedures specially to enhance the facial appearance and the requirement has expanded all over the world. This review was designed to identify and summarize the literature/evidence from different types of research/clinical studies conducted in facial resurfacing methodologies. A systematic review was designed and conducted in relation to facial resurfacing and rejuvenating techniques using data from PubMed, MEDLINE Cochrane database and Scopus. Systematic review protocol guidelines were used for research questions, aim of the review and PICO breakdown, eligibility criteria and used search engine and extraction of data. A minimum of 17 clinical studies and reviews were identified including randomized clinical trials. Significant results have been obtained in some clinical trials about techniques of rejuvenation and some studies have elucidated contrasting evidence. In most of the work, ablative and non-ablative laser therapy had been compared and we found that some results were compatible, and others showed contrasting evidence. In conclusion, more high-quality randomized control clinical trials are needed to decide on better safety and effectiveness of ablative or non-ablative laser therapy vs conventional treatments.

Index Terms- Laser Therapy, Face Resurfacing Technology

INTRODUCTION

Esthetic skin procedures have become a more demanded area among the younger generation of the world. There is a wide range of available techniques used for skin resurfacing [1]. Skin resurfacing treatment will mainly focus on the skin

complexion and removal of the top layer of the skin [1,2]. Our literature survey showed that this enhancing complexion procedure can be done by different methods [1-5]. In this review process, we concentrate on reviewing the therapeutic efficacy and safety of different types of laser technology, chemical peeling in rejuvenation and other methods which are also used in the practice. In all studies, objective of the facial resurfacing is mainly to remove of top layer of the skin by damaging the skin layer to regenerate the new young cells [3] All methods which are used currently, reduce fine and coarse wrinkles, fine lines, age specific spots and other blemishes [5,6]. Demand for rejuvenation is now more common among adult and aged generation in all over the world and had made their appearance of the face and neck drastically [6]. Some research findings showed that individual facial assessment is mandatory before the commencement of the therapy while some others have not given any data about the pretreatment evaluation. The safety, efficacy and post treatment complications have been compared and studied only in few studies [7,8].

CURRENT TREATMENT MODALITIES

Before the laser resurfacing, soft tissue fillers like fat, collagen or hyaluronic acid are commonly injected into wrinkled area in both neck and face [7,8,9]. With the practice of facial filling, research evidence showed that hyaluronic acid, calcium hydroxypaptite, fat is commonly used with safety [7,8]. Further, clinical trials showed that effective time duration of facial filling last for 6-12 months and repeated treatment therefore is required to regain the appearance when compared to the lase therapy [7,9,10]. Goldman *et al* with his research group had found out that there is no significant difference in the treatment groups of facial filling alone and facial filling followed with laser treatment on face [9]. In contrast, Fernando *et al* had

published a paper with a result of the concomitant use (same day) of laser and HA fillers for facial rejuvenation represents an effective and safe strategy which improves clinical results and patient's satisfaction [10]. This shows contradictory results a decision on the effectiveness and safety of combination filler/laser treatments [10]

CHEMICAL PEELING

Chemical peeling has been practiced on the face to remove the dead upper layer of the skin and stimulate the rejuvenation of the new skin cells [10]. Chemical peels are different types, and superficial and medium peels will improve the skin texture wrinkles and pigmentation and recovery will be smooth. Deep peels are used for more severe dermatological conditions and may result in swelling and blisters after the treatment [11]. Review published by Hesham *et al*, had identified that there is no difference in fractional erbium YAG with glycolic acid peel in facial treatment [11]. In support to this finding, Hassan *et al* also had conducted comparative study on CO₂ laser and chemical peeling and found out the high concentration of fractional CO₂ laser and chemical peeling had shown equally effective results in treatment of acne vulgaris [12]. But they specifically highlighted the fractional CO₂ laser is superior to mandelic acid chemical peeling in treatment of post acne scars even in high concentration [12]

MECHANISM OF LASER FACIAL RESURFACING

Different types of laser therapy is used to rejuvenate the skin in cosmetic industry to enhance the tone, texture and reduce pigmentation of the skin. Laser therapy uses the process of photothermolysis which will release a beam of light energy to the affected area of skin, and this will destroy the outer layer of the skin, which is epidermis. Laser beams cause heating of water at epidermis and vaporize the tissue to get filled by collagen fibers. The wavelength of the light must be chosen as per the color and the shape of the target area [3,5]. With this treatment, stimulation of dermis will be commenced with cellular and tissue repairing and collagen production. This revitalization procedure will make smooth and youthful appearance of the person [3-5].

As per the published reports in this study, we found that laser therapy is collectively used in removing a wide variety of scars in addition to the wrinkles [5]. Most of the studies identified that minimum recovery time is one of the advantages of laser treatment. Patients can resume day to day activities after the session and it has been recorded for the treatment of acne, birth marks, excessive hair growth problems, sagging skin conditions age related pigmentations and lesions, vascular lesion which are cosmetically affected, wrinkles and age spots and most importantly the precancerous lesions [3-6].

DIFFERENT TYPES OF LASER THERAPY

Laser therapy uses different types of lasers such as Erbium, CO₂ and diode lasers. There are two main types of ablative and non-ablative laser therapy.

Dermatologists and clinicians use fractional CO₂ lasers which are a mixture of 3 gases of 10-20% CO₂, 10-20% nitrogen and Helium. This is noninvasive and removes the damaged and aged skin and stimulates collagen formation and result in firm and healthy young skin. Erbium laser resurfacing has been done to remove surface level, deep lines and wrinkles. This has become popular due to the minimum burning of surrounding tissues [1,2,4]. If the skin is dark, treatment with Neodymium YAG has been considered because it has a longer wavelength that goes deeper into the skin effectively bypassing the melanin in the upper layer of the skin [2-4]

Many randomized control trials were reviewed to find out the best resurfacing technique [1-17]. Reviews articles were also studied and multiple authors in review articles had advocated improving the study designing and quality in future. [13,17]. In relation to the safety and efficacy, 1540-nm fractional erbium glass laser in the non-contact mode had been identified as safe and effective treatment for facial photodamage [15]. In addition to that another clinical trial showed that treatment of photodamage with a fractionated 1927-nm nonablative thulium laser is a promising new therapeutic option [16] and showed that two treatments with a 1927nm non-ablative fractionated thulium laser produced moderate to marked improvement in overall appearance and pigmentation with high patient satisfaction [16]. The response to treatment has been maintained for

three months follow up [16,17]. Another method of treatment for aging skin was identified as photo rejuvenation. Phototherapy had been very successful in controlling the regular pigmentation effectively. Light emitting diode (LED) is identified as novel light source and effective in wrinkles skin laxity. Yoon lee et al had conducted a prospective randomized double blind, split face clinical study and recommended that molecular level reactivation of skin rejuvenation with LED treatment and its safety [22,23].

A study conducted by Christopher *et al*, found out that techniques supported with artificial intelligence is much better than traditional methods of results of facial rejuvenating and resurfacing [24]. Review published by Macrene *et al* had suggested that multi-center clinical trial evidence for the better safety and efficacy with the fractional carbon dioxide laser treatment [25]. In contrast Yang *et al* researched that both nonablative fractional laser and ablative CO₂ fractional laser are effective and safe treatment for striae distensae of Asian skin. They further recommend that neither of treatment showed have not shown any greater clinical improvement than the other treatment. [26]

Considering all above research work and their evidence, treatment modalities for facial resurfacing had resulted different in individual work and vary with the type of the disease and the facial presentation. Article published in Journal of cosmetic and laser therapy by Fereshtech *et al* had identified that safety/efficacy and cost effectiveness of CO₂ fractional laser therapy in comparison to other methos of rejuvenation is high [18]. Cost for the treatment modalities had varied with the treatment type for the type of skin lesion. Farnoosh *et al* found out that the efficacy and safety of ablative laser were not higher than those of non-ablative laser in skin rejuvenation [19] Most importantly, minim invasion and permanent skin solution are the main reasons for the popularity of the laser skin treatment with very impressive results.

RECOMMENDATIONS

As per the finding with the former research and review articles, we only can recommend the performance of laser-based resurfacing is

considered as safe and efficient method of making expected skin appearance only with the hands of highly trained, knowledgeable professional.

Therefore, when you decide for laser skin resurfacing, with this study, we recommend discussing with the cosmetic surgeon o dermatologist about your focus on the treatment goals and the existing skill problem which you want to address and the expected results. In relation to efficacy and safety, considering all the above research work published, we identified contrasting results for recommending ablative laser and non-ablative laser therapy. Therefore, we need more high-quality randomized control clinical trials to decide on the safety and effectiveness of ablative or non-ablative laser therapy vs conventional treatments.

REFERENCES

1. Lipozencic J, Bukvic Mokos Z. Dermatologic lasers in the treatment of aging skin. *Acta Dermatovenerol Croat.* 2010;18(3):176–180. [PubMed] [Google Scholar]
2. Alster TS, Lupton JR. Lasers in dermatology. An overview of types and indications. *Am J Clin Dermatol.* 2001;2(5):291–303. [PubMed] [Google Scholar]
3. Smith KC, Schachter GD. YSGG 2790-nm Superficial Ablative and Fractional Ablative Laser Treatment. *Facial Plast Surg Clin North Am.* 2011;19(2):253–260. [PubMed] [Google Scholar]
4. Manstein D, Herron GH, Sink RK, et al. Fractional Photothermolysis: A New Concept for Cutaneous Remodeling Using Microscopic Patterns of Thermal Injury. *Lasers Surg Med.* 2004;34:426–438. [PubMed] [Google Scholar]
5. Malinowska S, Jaguś D, Woźniak W, Mlosek RK. Usefulness of high-frequency ultrasound in the monitoring of laser treatment of acne scars. *J Ultrason.* 2021;20(83):e279-e283. doi: 10.15557/JoU.2020.0049. Epub 2020 Dec 18. PMID: 33500795; PMCID: PMC7830061.
6. Alexiades-Armenakas MR, Dover JS, Arndt KA. Fractional laser skin resurfacing. *J Drugs*

- Dermatol.* 2012;11(11):1274–1287. [PubMed] [Google Scholar]
7. Foster KW, Moy RL, Fincher EF. Advances in plasma skin regeneration. *J Cosmet Dermatol.* 2008;7(3):169–179. [PubMed] [Google Scholar]
 8. Tierney EP, Hanke CW. Ablative fractionated CO₂ laser resurfacing for the neck: prospective study and review of the literature. *J Drugs Dermatol.* 2009 Aug;8(8):723-31. PMID: 19663109.
 9. Goldman MP, Alster TS, Weiss R. A randomized trial to determine the influence of laser therapy, monopolar radiofrequency treatment, and intense pulsed light therapy administered immediately after hyaluronic acid gel implantation. *Dermatol Surg.* 2007 May;33(5):535-42. doi: 10.1111/j.1524-4725.2007.33111.x. PMID: 17451575.
 10. Urdiales-Gálvez F, Martín-Sánchez S, Maíz-Jiménez M, Castellano-Miralla A, Lionetti-Leone L. Concomitant Use of Hyaluronic Acid and Laser in Facial Rejuvenation. *Aesthetic Plast Surg.* 2019 Aug;43(4):1061-1070. doi: 10.1007/s00266-019-01393-7. Epub 2019 May 9. Erratum in: *Aesthetic Plast Surg.* 2019 Sep 17;; PMID: 31073742; PMCID: PMC6742610.
 11. Hesham A. Shokeir 1 , Maha R. Abo Eittah 2 , Noura A. El Seessy 3 Comparative Study on Fractional Ablative Erbium: YAG Laser Versus Chemical Peeling in Treatment of Melasma Sys *Rev Pharm* 2020;11(12):593-599
 12. Hassan M. El-Fakahany, Walid M. Mohamed, and Sheyam S. Amer. Fractional CO₂ Laser and Chemical Peeling for Treatment of Acne and Acne Scars: A Comparative Study. *MJMR*, Vol. 31, No. 1, 2020, pages (66- 69).
 13. Halbina A, Trznadel-Grodzka E, Rotsztein H. Fractional laser therapy - the next step in alleviating the symptoms of skin aging (own observations). *Prz Menopauzalny.* 2014 May;13(2):132-5. doi: 10.5114/pm.2014.42716. Epub 2014 May 21. PMID: 26327843; PMCID: PMC4520352.
 14. Lapidoth M, Halachmi S, Cohen S, Amitai DB. Fractional CO₂ laser in the treatment of facial scars in children. *Lasers Med Sci.* 2014 Mar;29(2):855-7. doi: 10.1007/s10103-013-1305-6. Epub 2013 Mar 26. PMID: 23529372.
 15. Lapidoth M, Adatto M, Halachmi S. Treatment of actinic keratoses and photodamage with non-contact fractional 1540-nm laser quasi-ablation: an ex vivo and clinical evaluation. *Lasers Med Sci.* 2013 Feb;28(2):537-42. doi: 10.1007/s10103-012-1103-6. Epub 2012 Apr 27. PMID: 22538843.
 16. Weiss ET, Brauer JA, Anolik R, Reddy KK, Karen JK, Hale EK, Brightman LA, Bernstein L, Geronemus RG. 1927-nm fractional resurfacing of facial actinic keratoses: a promising new therapeutic option. *J Am Acad Dermatol.* 2013 Jan;68(1):98-102. doi: 10.1016/j.jaad.2012.05.033. Epub 2012 Oct 2. PMID: 23041112.
 17. Brauer JA, McDaniel DH, Bloom BS, Reddy KK, Bernstein LJ, Geronemus RG. Nonablative 1927 nm fractional resurfacing for the treatment of facial photopigmentation. *J Drugs Dermatol.* 2014 Nov;13(11):1317-22. PMID: 25607696.
 18. Pannucci CJ, Reavey PL, Kaweski S, Hamill JB, Hume KM, Wilkins EG, Pusic AL. A randomized controlled trial of skin care protocols for facial resurfacing: lessons learned from the Plastic Surgery Educational Foundation's Skin Products Assessment Research study. *Plast Reconstr Surg.* 2011 Mar;127(3):1334-1342. doi: 10.1097/PRS.0b013e318204361d. PMID: 21364435; PMCID: PMC3079206.
 19. Alexiades-Armenakas M, Sarnoff D, Gotkin R, Sadick N. Multi-center clinical study and review of fractional ablative CO₂ laser resurfacing for the treatment of rhytides, photoaging, scars and striae. *J Drugs Dermatol.* 2011 Apr;10(4):352-62. PMID: 21455544.
 20. Ansari, Fereshteh & Sadeghi-Ghyassi, Fatemeh & Yaaghoobian, Barmak. (2018). The

- clinical effectiveness and cost-effectiveness of fractional CO₂ laser in acne scars and skin rejuvenation: A meta-analysis and economic evaluation. *Journal of Cosmetic and Laser Therapy*. 20. 1-4. 10.1080/14764172.2017.1400173.
21. Seirafianpour, F., Pour Mohammad, A., Moradi, Y. *et al.* Systematic review and meta-analysis of randomized clinical trials comparing efficacy, safety, and satisfaction between ablative and non-ablative lasers in facial and hand rejuvenation/resurfacing. *Lasers Med Sci* 37, 2111–2122 (2022). <https://doi.org/10.1007/s10103-022-03516-0>
22. Lee SY, Park KH, Choi JW, Kwon JK, Lee DR, Shin MS, Lee JS, You CE, Park MY. A prospective, randomized, placebo-controlled, double-blinded, and split-face clinical study on LED phototherapy for skin rejuvenation: clinical, profilometric, histologic, ultrastructural, and biochemical evaluations and comparison of three different treatment settings. *J Photochem Photobiol B*. 2007 Jul 27;88(1):51-67. doi: 10.1016/j.jphotobiol.2007.04.008. Epub 2007 May 1. PMID: 17566756.
23. Ablon G. Phototherapy with Light Emitting Diodes: Treating a Broad Range of Medical and Aesthetic Conditions in Dermatology. *J Clin Aesthet Dermatol*. 2018 Feb;11(2):21-27. Epub 2018 Feb 1. PMID: 29552272; PMCID: PMC5843358.
24. Zachary T. Elliott, Alekya Bheemreddy, Michele Fiorella, Ann M. Martin, Vanessa Christopher, Howard Krein, Ryan Heffelfinger, Artificial intelligence for objectively measuring years regained after facial rejuvenation surgery, *American Journal of Otolaryngology*; 2023,44; 2.
25. Alexiades-Armenakas M, Sarnoff D, Gotkin R, Sadick N. Multi-center clinical study and review of fractional ablative CO₂ laser resurfacing for the treatment of rhytides, photoaging, scars and striae. *J Drugs Dermatol*. 2011 Apr;10(4):352-62. PMID: 21455544.
- 26, Yang YJ, Lee GY. Treatment of Striae Distensae with Nonablative Fractional Laser versus Ablative CO₂ Fractional Laser: A Randomized Controlled Trial. *Ann Dermatol*. 2011 Nov;23(4):481-9. doi: 10.5021/ad.2011.23.4.481. Epub 2011 Nov 3. PMID: 22148016; PMCID: PMC3229942.